

Office on Women's Health

Sarah Ehrhard Reid

Programs in OWH

SAFEVT

- Launching2025
- Supports survivors of violence in Missouri hospitals
- Opportunity for service connection

Maternal Mortality

- Pregnancy
 Associated
 Mortality
 Review
- Baby and Me Tobacco Free
- Awareness materials
- Notice of Pregnancy


Violence Prevention

- Shared risk and protective factors work
- Partnering
 with colleges,
 universities
 and the
 Missouri
 Coalition
 Against
 Domestic and
 Sexual
 Violence

Women's Health Initiatives

- Women's
 Health
 Advisory
 Committee
- Screening guides
- Extended Women's Health Services

SAFEVT Network






What is SAFEVT?

- SANE- Sexual assault nurse examiner
- SAFE- Sexual assault forensic exam
- •SAFEvT- Sexual assault forensic exam via telehealth



How does SAFEVT work?






What's can LPHAs do?

Support Survivors



Connect with victim service organizations



Work with hospitals



Support survivors' ongoing health needs



Pregnancy Associated Mortality Review 2018-2022

6

Leading causes of death. Mental health conditions, cardiovascular, infection, amniotic fluid embolism, homicide & hemorrhage.

82%

Deaths that occur after delivery up to one year postpartum.

40

The pregnancy-related mortality ratio per 100,000 live births for moms living in micropolitan counties.

2.5x

Pregnancy-related mortality ratio was 2.5 times higher for Black moms than white moms.

32.3


Pregnancy-related mortality ratio per 100.000 live births.

Key Findings 2018-2022


4 in 5

Deaths determined to be preventable (80%).

Underlying Causes of Pregnancy-Related Deaths,



Timing of Pregnancy-Related Deaths, 2018-2022



Ordering Resources or Connecting with OWH

womenshealth@health.mo.gov